

What is claimed is:

1. A composition of matter comprising: a biomolecule in combination with a metal particle, wherein said metal particle and said biomolecule are positioned at a distance apart sufficient to adjust intrinsic emission of electromagnetic radiation from the biomolecule in response to an amount of exciting electromagnetic radiation.

2. The composition of claim 1, wherein the biomolecule comprises a nucleic acid.

3. The composition of claim 1, wherein the biomolecule comprises a purine or pyrimidine.

4. The composition of claim 1, wherein the biomolecule comprises a nucleoside or nucleotide.

5. The composition of claim 1, wherein the biomolecule comprises an oligonucleotide.

6. The composition of claim 1, wherein the biomolecule comprises a protein.

7. The composition of claim 1, wherein the biomolecule comprises an amino acid.

8. The composition of claim 1, wherein the biomolecule comprises a lipid.

9. The composition of claim 1, wherein the biomolecule comprises a sugar moiety.

10. The composition of claim 1, wherein the metal particle is at a distance of about 50 Å to about 2000 Å from the biomolecule.

11. The composition of claim 1, wherein the metal particle comprises a noble metal.

12. The composition of claim 11, wherein the noble metal is selected from the group consisting of rhenium, ruthenium, rhodium, palladium, silver, osmium, iridium, platinum, and gold.

13. The composition of claim 1, wherein the metal particle is sub-wavelength in size.

14. The composition of claim 1, wherein the biomolecule is linked to the metal particle.

15. A method for increasing the intrinsic fluorescence of a biomolecule, said method comprising the step of:

positioning a metal particle and said biomolecule at a distance apart sufficient to increase
the electromagnetic emission from said biomolecule in response to an amount of exciting
radiation.

16. A method for detecting a biomolecule, said method comprising the steps of:

(a) positioning a metal particle and a biomolecule at a distance apart sufficient to
manipulate the electromagnetic emission from said biomolecule;

(b) exposing said biomolecule to an amount of exciting radiation; and

(c) detecting the electromagnetic emission from said biomolecule.

17. A method for manipulating fluorescence intensity of a biomolecule, said method comprising the steps of:

(a) increasing the rate of radiative decay of the biomolecule by positioning the
biomolecule adjacent to a metal particle; and

(b) exposing the biomolecule to an amount of exciting radiation.

18. A method for detecting the presence of a nucleic acid sequence in a sample, said method comprising the steps of:

(a) providing a sample;

(b) adding a nucleic acid sequence linked to a metal particle;

(c) exposing the sample to an amount of exciting radiation;

(d) detecting the fluorescence; and

(e) determining the presence of the nucleic acid sequence based on the detection of the fluorescence.

19. A method for increasing the fluorescence intensity of a fluorescently labeled biomolecule, said method comprising the steps of:

5

(a) labeling a biomolecule with a fluorophore;

(b) positioning the labeled biomolecule next to a metallic particle such that in response to an amount of exciting radiation, the fluorophore emits radiation.

20. A method for increasing fluorescence energy transfer on a fluorescently labeled biomolecule, said method comprising the steps of:

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

(a) labeling a biomolecule with a donor fluorophore and an acceptor fluorophore;

(b) positioning the labeled biomolecule adjacent to a metal particle such that in response to an amount of exciting radiation, the donor fluorophore transfers energy to the acceptor fluorophore causing the acceptor fluorophore to emit electromagnetic radiation.

21. A method for increasing fluorescent intensity of a fluorophore, said method comprising the steps of:

(a) positioning a fluorophore adjacent to a metal particle; and

(b) exciting said fluorophore with a plurality of photons.

22. A method for increasing fluorescent intensity of a biomolecule, said method comprising the steps of:

20

(a) positioning a biomolecule adjacent to a metal particle; and

(b) exciting said biomolecule with a plurality of photons.

23. A method for selectively enhancing the region of electromagnetic emission of a sample, said method comprising the steps of:

- (a) directing a metal particle to a region of interest in the sample; and
- (b) providing an amount of exciting radiation in the region of interest.

24. A method for selectively enhancing the region of electromagnetic emission of a sample, said method comprising the steps of:

- (a) directing a metal particle to a region of interest in the sample;
- (b) contacting the sample with a fluorophore;
- (c) exposing the sample to an amount of exciting radiation.

25. A method for increasing fluorescence energy transfer on a fluorescently labeled biomolecule, said method comprising the steps of:

- (a) labeling a first biomolecule with a donor fluorophore
- (b) labeling a second biomolecule with an acceptor fluorophore;
- (c) positioning the labeled biomolecules adjacent to a metal particle such that in response to an amount of exciting radiation, the donor fluorophore transfers energy to the acceptor fluorophore increasing the emission of electromagnetic radiation of the acceptor fluorophore.

26. A microarray system comprising:

a solid support, wherein the solid support is coated with metal particles; and

a matrix having an array of biomolecules attached to the support such that when a labeled probe hybridizes to the biomolecules, the fluorescence of the labeled probe increases in response to an amount of exciting radiation.

27. A composition of matter comprising: a biomolecule in combination with a metal surface, wherein said metal surface and said biomolecule are positioned at a distance apart sufficient to adjust intrinsic emission of electromagnetic radiation from the biomolecule in response to an amount of exciting electromagnetic radiation.